

FINGERS OF GOD

A CRITIQUE OF REES'S THEORY OF PRIMORDIAL GRAVITATIONAL RADIATION

J.C. Jackson¹Royal Greenwich Observatory
Herstmonceux Castle
Sussex
UK

(Received 1971 November 12)

SUMMARY

Very long wavelength universal gravitational waves cannot now produce in clusters of galaxies velocity dispersions greater than that which these systems would possess if they were expanding with the Universe, if the Universe is not younger than 10^{10} yr and Hubble's constant is not less than 50 km/sec/ Mpc. A diagram shows that actual velocity dispersions are significantly greater than this limit.

¹Present address: Division of Mathematics, School of Computing, Engineering and Information Sciences, Northumbria University, Newcastle NE1 8ST, UK

INTRODUCTION

The persistently negative results of searches for enough ordinary matter in clusters of galaxies to account for their large velocity dispersions will probably stimulate interest in ‘radical’ solutions of this problem. I shall discuss here one such proposal, namely Rees’s hypothesis (1971) that these systems are interacting with very long wavelength gravitational waves. Before doing so, however, I shall present a diagram which shows very clearly what I consider to be the essential feature of this phenomenon; it is this feature which cannot be explained by Rees’s theory in its present form.

Fig. 1 shows the spatial distribution of all galaxies with right ascensions in the range 10-11 hr, and known positive radial velocities (relative to the local group) less than 2000 km s^{-1} . The ‘distance’ D used in this plot is defined as $\text{velocity}/100 \text{ km/sec/Mpc}$. The galaxies appear to fall into long chains or cigar-shaped configurations, all pointing at the Earth. Unless one is prepared to assign to the Earth a very special place in the Universe, one must conclude that D is not a good distance indicator, and that in reality the galaxies exist in roughly spherical configurations whose internal velocity dispersions are several times that which would be observed if these systems were expanding with the Universe. (I shall call the latter the internal Hubble velocity.) It is this aspect of these small groups, rather than their mass discrepancy, which is odd, as in many cases the virial theorem velocity corresponding to the visible mass is considerably smaller than the internal Hubble velocity. I shall show that gravitational waves could at most produce a velocity dispersion equal to the internal Hubble velocity.

The groups apparent in Fig. 1 coincide with members of de Vaucouleurs’ complete list (1971) of small groups within 17 Mpc of the local group, which was compiled using distance indicators other than velocity. An analysis of this list gives a mean value of the ratio internal velocity dispersion/internal Hubble velocity ~ 5 . Although the precise value of this number is affected by uncertainties in group membership, diagrams such as Fig. 1 convince one that the groups are real and that this ratio is significantly greater than unity.

The apparent instability exhibited by de Vaucouleurs’ groups is not particularly violent; their crossing times are longer than 10^9 yr (Rood, Rothman & Turnrose 1970). Systems with much shorter crossing times are known (see for example Burbidge & Sargent 1970); however, following Rees I shall restrict my discussion to groups with long crossing times.

GRAVITATIONAL WAVES

The interaction of a plane gravitational wave with a cluster of galaxies can be solved exactly in the linearized version of general relativity (for a suitable theoretical background see Weber 1961). Such a wave, moving along the x -axis with angular frequency ω , is described by the metric

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$$

where $\eta_{\mu\nu}$ is the Minkowski metric, and the first order perturbation $h_{\mu\nu}$ is given by

$$h_{\mu\nu} = A e_{\mu\nu} \exp \left[i\omega \left(t - \frac{x}{c} \right) \right]$$

($\mu, \nu = 0, 1, 2, 3$ indicates t, x, y, z components respectively).

A is the amplitude of the wave, and for a fully polarized one the polarization tensor $e_{\mu\nu}$ can be




Figure 1: The apparent space distribution of nearby galaxies with $10h \leq RA \leq 11h$, assuming radial velocity to be a good distance indicator.

written as

$$e_{\mu\nu} = \frac{1}{\sqrt{2}}(\delta_{\mu}^2 \delta_{\nu}^2 - \delta_{\mu}^3 \delta_{\nu}^3)$$

where δ_{ρ}^{σ} is the Kronecker δ symbol. Consider now the interaction of this radiation with a system of N point particles, with masses m_i , and position vectors (x_i, y_i, z_i) and velocities (u_i, v_i, w_i) in centre of mass coordinates; the gravitational interaction between these particles will be neglected. (The masses m_i could be omitted from this discussion; the advantage of including them is that the resulting mass weighted quantities are readily available for many systems.) Their accelerations relative to the centre of mass are related to the Riemann tensor $R_{\mu\nu\rho\sigma}$ via the equation of geodesic deviation:

$$\begin{aligned}\ddot{x}_i &= 0 \\ \ddot{y}_i &= R_{0202}y_i = -\frac{1}{2}\ddot{h}_{22}y_i = \frac{1}{2}\omega^2 A e_{22}y_i \exp\left[i\omega\left(t - \frac{x}{c}\right)\right] \\ \ddot{z}_i &= R_{0303}z_i = -\frac{1}{2}\ddot{h}_{33}z_i = \frac{1}{2}\omega^2 A e_{33}z_i \exp\left[i\omega\left(t - \frac{x}{c}\right)\right]\end{aligned}$$

where a dot indicates a time derivative. If, as in the situation envisaged by Rees, the particles do not suffer any sensible change of position during the course of an oscillation, and the only appreciable velocities are those induced by the wave, these equations integrate to

$$u_i = 0$$

$$v_i = \frac{1}{2}\omega A e_{22} y_i \exp \left[i\omega \left(t - \frac{x}{c} \right) - i\pi/2 \right]$$

$$w_i = \frac{1}{2}\omega A e_{33} z_i \exp \left[i\omega \left(t - \frac{x}{c} \right) - i\pi/2 \right]$$

Let M be the total mass of the system, and define the velocity dispersion V by

$$MV^2 = \text{time average of } \left[\sum_i m_i (|u_i|^2 + |v_i|^2 + |w_i|^2) \right].$$

Then

$$MV^2 = \frac{1}{16}\omega^2 A^2 \sum_i m_i (y_i^2 + z_i^2) \sim \frac{1}{16}\omega^2 A^2 \times \frac{2}{3} \sum_i m_i (x_i^2 + y_i^2 + z_i^2)$$

so that

$$V^2 \sim \frac{1}{24}\omega^2 A^2 R_I^2$$

where R_I is the radius of gyration of the system about its centre of mass:

$$MR_I^2 = \sum_i m_i (x_i^2 + y_i^2 + z_i^2).$$

We are concerned with wavelengths $\lambda \ll$ radius of the Universe, and the effective mass density of this radiation is (Isaacson 1968)

$$\rho_G = \frac{A^2 \omega^2}{64\pi G}.$$

Thus in terms of ρ_G , we have

$$\frac{V}{R_I} = \left(\frac{8}{3}\pi G \rho_G \right)^{1/2}. \quad (1)$$

For a truly plane wave, this expression is valid for all values of λ ; Rees notes that it would be unreasonable to suppose that the planeness extends over regions much greater than λ , so that if $\lambda < R_I$ this expression should be reduced by a factor λ/R_I . However, when $\lambda \ll R_I$, the gravitational waves will act as a smoothed out medium of density ρ_G for the cluster as well as for the Universe, and the velocity dispersion (1) is precisely that which the ordinary gravity of such a medium (if static) would produce in the system. Thus equation (1) is possibly valid for all values of λ , although the last point is not clear as the medium would be expanding on a timescale comparable with the time taken by members to cross the system. Equation (1) certainly gives an upper limit for Rees's mechanism whatever the value of λ . Note that the wave has no associated gravitational redshift to augment the effect, as $h_{00} = 0$.

The results of this calculation agree with Rees' order of magnitude estimates; however, it is important to establish that R_I is the correct radius to use in this context, as radii defined in other

ways differ considerably in magnitude. For example, the radius R_Ω , defined so that $-GM^2/R_\Omega$ is the gravitational potential energy of the system, appears in applications of the virial theorem to groups (see for example Rood, Rothman & Turnrose 1970), and is typically larger than R_I by a factor of 3. The use of R_Ω in equation (1) would therefore considerably overestimate the effect. V is maximized by maximizing ρ_G , i.e. by considering a universe in which almost all the mass-energy is in the form of radiation. The observed deceleration of the Universe would then be due entirely to this radiation, and the deceleration parameter q_0 would be given by (Sandage 1961)

$$q_0 = \frac{8}{3}\pi G\rho_G H_0^{-2}$$

where H_0 is Hubble's constant. In this case, equation (1) becomes

$$\frac{V}{R_I} = q_0^{1/2} H_0. \quad (2)$$

It is easy to show that under these conditions q_0 is almost certainly less than unity. Thus we see that gravitational waves could at most produce a velocity dispersion in a system equal to that which would be produced by the Hubble expansion. If the observed values of V/R_I really were of order H_0 , rather than $5H_0$, it is doubtful that astronomers would consider that clusters of galaxies presented any problems. The upper limit on q_0 is derived as follows; let t_0 be the present age of the Universe; then the inequalities

$$t_0 \geq 10^{10} \text{ yr} \quad H_0 \geq 50 \text{ km/sec/Mpc}$$

are almost certainly true. Thus

$$H_0 t_0 \geq \frac{1}{2} \quad (3)$$

which, for a universe filled with radiation, implies

$$q_0 \leq 1.$$

(The value $q_0 = 25$ required to produce the observed values of V/R_I would need $H_0 \leq 17$ km/sec/Mpc if $t_0 \geq 10^{10}$ yr.)

I shall conclude by noting that another recently suggested 'radical' solution of this problem (Jackson 1970; see also Forman 1970; Paal 1964), which attributes the velocity dispersions to a negative cosmological constant, suffers from similar, but less severe, problems. In fact this theory also gives rise to equation (2); however, in the appropriate universe, inequality (3) leads to

$$q_0 \leq 5.4$$

and thus

$$\frac{V}{R_I} < 2.3.$$

ACKNOWLEDGMENTS

It is a pleasure to thank Dr D. Lynden-Bell for several stimulating conversations on this subject.

Royal Greenwich Observatory, Herstmonceux Castle, Sussex

REFERENCES

- Burbidge, E. M. & Sargent, W. L. W., 1971. Pont. Acad. Scient. Scripta Varia 35 (Vatican Conference on the Nuclei of Galaxies).
Forman, W. R., 1970. Astrophys. J., 159, 719.
Isaacson, R. A., 1968. Phys. Rev., 166, 1272.
Jackson, J. C., 1970. Mon. Not. R. astr. Soc., 148, 249.
Paal, G., 1964. Acts. Phys. Hung., 17, 379.
Rood, H. J., Rothman, V. C. A. & Turnrose, B. E., 1970. Astrophys. J., 162, 411.
Rees, M. J., 1971. Mon. Not. R. astr. Soc., 154, 187.
Sandage, A., 1961. Astrophys. J., 133, 355.
Vaucouleurs, G. de, 1971. Stars and Stellar Systems, Vol. 9, University of Chicago Press, in press.
Weber, J., 1961. General Relativity and Gravitational Waves, Interscience, New York.